



## **Drought & Lessons Learned**

**Jeanine Jones, California Department of Water Resources**

# First a Word About Defining Drought

## When Does “Dry” Become “Drought”?


- Meteorological drought
- Hydrological drought
- Regulatory drought
- Drought indices, US Drought Monitor
- Sector-based definitions
- **Drought is a function of impacts (which are typically regional or local)**

# When Does “Drought” Become “Drought Emergency”?

- Depends on impacts, and ability to mitigate impacts
- Drought differs from traditional “emergencies” (flood, fire, etc) in its very slow timescale
- California Emergency Services Act
  - Role of local government (counties)
  - Role of state




# Variability of Western Precipitation





# Reconstructed Streamflows & Drought Periods



## USING TREE-RINGS TO RECONSTRUCT STREAMFLOW

A tree-ring reconstruction is a set of tree-ring width data that have been calibrated with an instrumental or gaged record of a hydrologic or climatic variable such as annual streamflow or precipitation. The reconstruction, based on a statistical model that describes the relationship between tree growth and the gage record, extends that record back hundreds of years into the past.

Tree growth in dry climates is limited by water availability. Trees that provide the best information about hydroclimatic variability are those particularly sensitive to variations in moisture. These include species such as blue oak, ponderosa pine, Douglas fir, and western juniper, usually growing at lower elevations in sparse stands on dry and rocky sites where soil moisture storage is minimal.

Tree-ring reconstructions of hydroclimatic variables are developed from tree-ring chronologies. A tree-ring chronology is a time-series of annual values derived from the ring-width measurements of 10 or more trees of the same species at a single site. To create a tree-ring chronology, cores from the sampled trees at each site are cross-dated (i.e. patterns of narrow and wide rings are matched from tree to tree) to account for missing or false rings, so that every annual ring is absolutely dated to the correct year. Then all rings are measured to the nearest thousandth of a millimeter using a computer-assisted measuring device. After growth-related trends unrelated to climate are statistically removed, the ring width values from all sampled trees for each year are averaged to create a time series of annual ring width indices. The complete series of ring width indices from a site is called a tree-ring chronology.

Once a gaged record of interest is selected for reconstruction, a set of tree-ring chronologies from the region near the gage is calibrated with the gage record to form a reconstruction model. A statistical technique called multiple linear regression is commonly used. The reconstructions are evaluated by comparing the observed gage values with the reconstructed values by assessing the amount of variance in the gage record that is explained by the reconstruction.

**DROUGHTS PRIOR TO THE HISTORICAL RECORD**


The period of reliably measured streamflows for rivers throughout the West seldom reaches beyond 100 years, which represents only a fraction of climatologically modern time. As these streamflow reconstructions show, there have been droughts prior to the historical period that were more severe - particularly in duration - than those in the measured record. The reconstructed record captures a broader range of hydrologic variability than does the historical record, making reconstructions useful for drought preparedness planning. Of particular interest from a scientific perspective is the Medieval Climate Anomaly, a time during which sustained severe drought gripped much of the western United States, as exemplified illustrated in the Sacramento, San Joaquin, and Colorado River reconstructions.



Data source: Work performed by the University of Arizona under contract to the California Department of Water Resources. CDWR Agreements 460000382 (David Meko, 2006) and 4600008859 (David Meko, Connie Woodhouse, Ramo Touchan, 2014)




# Southern California Local Sources Example





# Annual California Runoff



# California's 20<sup>th</sup> & 21<sup>st</sup> Century Statewide Droughts (consecutive dry years)

- 1918-20
- 1922-24
- 1929-34
- 1947-50
- 1959-61

- 1976-77
- 1987-92
- 2007-09
- 2012-2016
- 2020- ??



# 21<sup>st</sup> Century & 20<sup>th</sup> Century Droughts Not the Same



# 2012-16

- Included warmest years on record, record low statewide snowpack
- State response actions not seen since 1976-77
- First-ever zero CVP ag contractor allocations
- About 500,000 acres fallowed
- First-ever state emergency response for areas of dry private residential wells
- First-ever use of InSAR to monitor statewide land subsidence

# California's Present Drought

- Zero allocation to most CVP ag contractors in WY 2021 and 2022, CVP M&I health & safety allocation in WY 2022, 5% SWP allocation
- 2022 large-scale urban water use restrictions in Southern California due to infrastructure limitations
- First Lower Colorado River Basin shortage pursuant to the Interim Guidelines
- Record low Lake Oroville elevation in 2021, Hyatt PP unable to generate
- 70% statewide snowpack in WY 2021, yet runoff comparable to 2014-2015
- Groundwater impacts similar to San Joaquin Valley in 2012-16 now seen in parts of Sacramento Valley



**Historically the Colorado River has been California's most reliable drought supply**



# What Has Changed Over Decades?

- Extensive interconnections now among largest water projects & urban suppliers
- Much greater experience with water transfers
- New groundwater management legislation
- Beginning in mid-1990s, substantial state grant funding for local projects
- It's getting warmer
- Wildfire risk increasing
- Increased acreage of permanent plantings
- Land subsidence in historically unaffected areas
- Small water system/private well owner problems becoming more widespread

# More Recent Changes

## Old

- Multi-year drought normal in reconstructed paleo & historical records
- Severely reduced CVP & SWP allocations
- Groundwater overdraft & land subsidence
- Impacts in San Joaquin Valley

## New

- Droughts occur in warming climate, exacerbates impacts
- First Lower Colorado River Basin shortage declared, SWP & CVP health & safety allocations
- Early stages of SGMA implementation
- Impacts in Sac Valley




# Key Drought Challenges, California's Progress Since 2007-09

- Statewide coverage of groundwater level data in major aquifers ✓
  - CASGEM legislation in 2009, SGMA legislation in 2014
- Small water systems ✓
  - Multiple legislative provisions for mandatory system consolidation beginning in 2016, long-term resilience funding legislation in 2019 (\$130M annually for 10 years), DWR drought response grants in 2021 & 2022
- Seasonal precipitation forecasting X
  - No improvement (requires action by NOAA)

# Historical Skill of NOAA Seasonal Outlooks – Not Usable for Water Management

Seasonal (Lead 0.5 Months) Precipitation Heidke Skill Score  
DJF Manual Forecasts From 1995 to 2022



# Catastrophic Wildfire Risk

- 1991 Oakland Hills fire (25 lives lost)
- October – November 2003 Southern California wildfires (22 lives lost)
- October 2007 Southern California wildfires (1 million people evacuated)
- 2017 Tubbs Fire, 2018 Camp Fire, 2021 Dixie Fire (urban water distribution system destruction)
- All but 2 of the state's 20 largest & 20 most damaging fires have occurred from 2000 onward





# Precautionary Evacuations at Hyatt & Keswick Powerplants



# Wildfire Damage to Water Infrastructure



# **Lessons Learned from Past Droughts**



JANUARY 2020

# California's Most Significant Droughts:

Comparing  
Historical and  
Recent  
Conditions

## Report to the Legislature on the 2012-2016 Drought

As Required by Chapter 340 of 2016

March 2021

# Lessons Learned From Past Droughts

- Impacts are highly site-specific, and vary depending on the ability of water users to invest in reliability
- Small water systems on fractured rock groundwater sources are most at risk of public health and safety impacts
- Larger urban water agencies can manage multiple years of drought with minimal impacts to their customers

# Expected Impacts of Multi-Year Drought

- **Unmanaged systems**
  - **Risk of catastrophic wildfire** (health & safety, economic)
  - Non-irrigated agriculture (livestock grazing)
  - Fish & wildlife (e.g., salmonids)
- **Managed systems**
  - **Small water systems** (health & safety)
  - Irrigated agriculture
  - Green industry (urban water supplies)
  - Fish & wildlife (e.g., wildlife refuges, salmonids)
  - Other environmental (e.g., land subsidence)



# Lessons Learned from Recent Droughts

- Act sooner when dry conditions emerge
- Recognize that increased temperatures are creating new or intensified impacts
- Plan for cutbacks in historical irrigation deliveries affecting shallow drinking water wells due to absence of groundwater recharge sources or compensatory construction of deeper irrigation wells
- Plan for wildfire impacts
- Transition from thinking of drought as an occasional emergency to thinking in terms of creating resiliency in a more arid climate



# Different State Drought Response Framework in Each Major Drought

- Reflecting differing conditions at the time
- Statewide drought emergency proclamation used in 3 of the “Big 5” droughts, but different type of use in each of those 3 cases
- Substantial state financial assistance response in only 2 of the Big 5 droughts
- **Cannot assume that substantial state financial available in future droughts, depends on state budget circumstances**

# State Capacity Development for Drought Response

- 1987-92: water transfers
- 2012-16: bond funding for initial groundwater data improvements
- 2020-??: recognition of role played by climate change
- Capacity-building investments needed for climate change response tools (forecasting & planning)

